FANCA gene

Fanconi anemia complementation group A

Normal Function

The FANCA gene provides instructions for making a protein that is involved in a cell process known as the Fanconi anemia (FA) pathway. The FA pathway is turned on (activated) when the process of making new copies of DNA, called DNA replication, is blocked due to DNA damage. The FA pathway is particularly responsive to a certain type of DNA damage known as interstrand cross-links (ICLs). ICLs occur when two DNA building blocks (nucleotides) on opposite strands of DNA are abnormally attached or linked together, which stops the process of DNA replication. ICLs can be caused by a buildup of toxic substances produced in the body or by treatment with certain cancer therapy drugs.

The FANCA protein is one of a group of proteins known as the FA core complex. The FA core complex is composed of eight FA proteins (including FANCA) and two proteins called Fanconi anemia-associated proteins (FAAPs). This complex activates two proteins, called FANCD2 and FANCI, by attaching a single molecule called ubiquitin to each of them (a process called monoubiquitination). The activation of these two proteins, which attach (bind) together to form the ID protein complex, attract DNA repair proteins to the area of DNA damage so the error can be corrected and DNA replication can continue.

Health Conditions Related to Genetic Changes

Fanconi anemia

More than 450 mutations in the *FANCA* gene have been found to cause Fanconi anemia, a disorder characterized by a decrease in bone marrow function, an increased cancer risk, and physical abnormalities. Mutations in the *FANCA* gene are responsible for 60 to 70 percent of all cases of Fanconi anemia. These mutations change single DNA building blocks (nucleotides) or insert or delete pieces of DNA in the *FANCA* gene. Some mutations allow production of a FANCA protein that has some residual function; other mutations prevent the production of any FANCA protein. Mutations that prevent all protein production usually lead to a shortage of blood cells at an earlier age and increase the risk of developing cancer of the blood-forming cells (leukemia) as compared to mutations that allow for some FANCA protein production.


Mutations in the *FANCA* gene lead to a nonfunctional FA core complex, which disrupts the entire FA pathway. As a result, DNA damage is not repaired efficiently

and ICLs build up over time. The ICLs stall DNA replication, ultimately resulting in either abnormal cell death due to an inability make new DNA molecules or uncontrolled cell growth due to a lack of DNA repair processes. Cells that divide quickly, such as bone marrow cells and cells of the developing fetus, are particularly affected. The death of these cells results in the decrease in blood cells and the physical abnormalities characteristic of Fanconi anemia. When the buildup of errors in DNA leads to uncontrolled cell growth, affected individuals can develop leukemia or other cancers.

Chromosomal Location

Cytogenetic Location: 16q24.3, which is the long (q) arm of chromosome 16 at position 24.3

Molecular Location: base pairs 89,737,551 to 89,816,658 on chromosome 16 (Homo sapiens Annotation Release 108, GRCh38.p7) (NCBI)



Credit: Genome Decoration Page/NCBI

Other Names for This Gene

- FA
- FAA
- FACA
- FANCA HUMAN
- Fanconi anemia, complementation group A

Additional Information & Resources

Educational Resources

- Madame Curie Bioscience Database: FANCA https://www.ncbi.nlm.nih.gov/books/NBK6302/#A45460
- Madame Curie Bioscience Database: The FA "Nuclear Core Complex" https://www.ncbi.nlm.nih.gov/books/NBK6302/#A45480

GeneReviews

 Fanconi Anemia https://www.ncbi.nlm.nih.gov/books/NBK1401

Scientific Articles on PubMed

PubMed

https://www.ncbi.nlm.nih.gov/pubmed?term=%28FANCA%5BTIAB%5D%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+1440+days%22%5Bdp%5D

OMIM

 FANCA GENE http://omim.org/entry/607139

Research Resources

- Atlas of Genetics and Cytogenetics in Oncology and Haematology http://atlasgeneticsoncology.org/Genes/FA1ID102.html
- ClinVar https://www.ncbi.nlm.nih.gov/clinvar?term=FANCA%5Bgene%5D
- HGNC Gene Family: Fanconi anemia complementation groups http://www.genenames.org/cgi-bin/genefamilies/set/548
- HGNC Gene Symbol Report http://www.genenames.org/cgi-bin/gene_symbol_report?q=data/ hgnc_data.php&hgnc_id=3582
- NCBI Gene https://www.ncbi.nlm.nih.gov/gene/2175
- The Rockefeller University: Fanconi Anemia Mutation Database: FANCA http://www2.rockefeller.edu/fanconi/genes/jumpa
- UniProt http://www.uniprot.org/uniprot/O15360

Sources for This Summary

- Deakyne JS, Mazin AV. Fanconi anemia: at the crossroads of DNA repair. Biochemistry (Mosc).
 2011 Jan;76(1):36-48. Review.
 Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/21568838
- OMIM: FANCA GENE http://omim.org/entry/607139

- Kee Y, D'Andrea AD. Expanded roles of the Fanconi anemia pathway in preserving genomic stability. Genes Dev. 2010 Aug 15;24(16):1680-94. doi: 10.1101/gad.1955310. Review. Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/20713514
 Free article on PubMed Central: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2922498/
- Taniguchi T, D'Andrea AD. Molecular pathogenesis of Fanconi anemia: recent progress. Blood. 2006 Jun 1;107(11):4223-33. Epub 2006 Feb 21. Review.
 Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/16493006
- de Winter JP, Joenje H. The genetic and molecular basis of Fanconi anemia. Mutat Res. 2009 Jul 31;668(1-2):11-9. doi: 10.1016/j.mrfmmm.2008.11.004. Epub 2008 Nov 14. Review.
 Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/19061902

Reprinted from Genetics Home Reference: https://ghr.nlm.nih.gov/gene/FANCA

Reviewed: January 2012 Published: March 21, 2017

Lister Hill National Center for Biomedical Communications U.S. National Library of Medicine National Institutes of Health Department of Health & Human Services